

Cambridge IGCSE[®]

CANDIDATE NAME			
CENTRE NUMBER		CANDIDATE NUMBER	
CHEMISTRY			0620/04
Paper 4 Theory	y (Extended)	For ex	camination from 2020
SPECIMEN PA	PER		

Candidates answer on the question paper.

No additional materials are required.

READ THESE INSTRUCTIONS FIRST

Write your centre number, candidate number and name on all the work you hand in. Write in dark blue or black pen. You may use an HB pencil for any diagrams, graphs or rough working. Do not use staples, paper clips, glue or correction fluid. DO **NOT** WRITE IN ANY BARCODES.

Answer **all** questions.

Electronic calculators may be used. You may lose marks if you do not show your working or if you do not use appropriate units. A copy of the Periodic Table is printed on page 20.

At the end of the examination, fasten all your work securely together. The number of marks is given in brackets [] at the end of each question or part question.

This document consists of 18 printed pages and 2 blank pages.

1 hour 15 minutes

substance	melting point / °C	boiling point / °C	electrical conductivity as a solid	electrical conductivity as a liquid
Α	839	1484	good	good
В	-188	-42	poor	poor
С	776	1497	poor	good
D	-117	78	poor	poor
Е	1607	2227	poor	poor
F	-5	102	poor	good
a) Which sul	bstance could b	e a metal?		

The following table gives information about six substances. 1

[1] (b) State all the substances that are liquid at room temperature. [1] (c) Which substance could have a macromolecular structure similar to that of silicon(IV) oxide? [1] (d) Which substance could be propane? [1] (e) Which substance could be sodium chloride? [1] [Total: 5] **2** The table gives the composition of three particles.

particle	number of protons	number of electrons	number of neutrons
Α	15	15	16
В	15	18	16
С	15	15	17

- (a) What is the evidence in the table for each of the following?
 - (i) Particle **A** is an atom.

 [1]
 (ii) A, B and C are all particles of the same element.

 [1]
 (iii) Particles A and C are isotopes of the same element.

 [1]
 (iii) Particles A and C are isotopes of the same element.

 [2]
 (b) (i) What is the electronic structure of particle A?

 [1]
 (ii) Is element A, a metal or a non-metal? Give a reason for your choice.

 [1]
 [1]

 [1]
 [1]

 [1]
 [1]

- **3** Kinetic theory explains the properties of matter in terms of the arrangement and movement of particles.
 - (a) Nitrogen is a gas at room temperature. Nitrogen molecules, N₂, are spread far apart and move in a random manner at high speed.
 - (i) Draw the electronic structure of a nitrogen molecule. Show only the outer electron shells.

[2]


(ii) Compare the movement and arrangement of the molecules in solid nitrogen to those in nitrogen gas.

[3]

(b) A sealed container contains nitrogen gas. The pressure of the gas is due to the molecules of the gas hitting the walls of the container. Use the kinetic theory to explain why the pressure inside the container increases when the temperature is increased.

[2]

The following apparatus can be used to measure the rate of diffusion of a gas.




The following results were obtained.

gas	temperature /°C	rate of diffusion in cm ³ /min
nitrogen	25	1.00
chlorine	25	0.63
nitrogen	50	1.05

(c) (i) Explain why nitrogen gas diffuses faster than chlorine gas.

		[2]
(ii)	Explain why the nitrogen gas diffuses faster at the higher temperature.	
		[1]
	[Total:	10]

- (a) (i) State two differences in the physical properties of chromium and sodium.
 [2]
 (ii) State two differences in the chemical properties of chromium and sodium.
 [2]
 - (b) Chromium is used to electroplate steel objects. The diagram shows how this could be done.



(ii)	The formula of the chromium(III) ion is Cr^{3+} and of the sulfate ion is SO_4^{2-} . Give the formula of chromium(III) sulfate.	[2] he
		[1]
(iii)		
	[[2]
(iv)	A colourless gas, which relights a glowing splint, is formed at the positive electrod (anode).	de
	State the name of this gas.	
		[1]


Chromium is a transition element.

(v) During electroplating, it is necessary to add more chromium(III) sulfate but during copper plating using a copper anode, it is not necessary to add more copper(II) sulfate.

Explain this difference.

..... [2] [Total: 12]

5 Iron is extracted from its ore, hematite, in the blast furnace.



Describe the reactions involved in this extraction.

Include one equation for a redox reaction and one for an acid/base reaction.

••••••	
••••••	
••••••	
	[5
	[0]
	[Total: 5
	• •

BLANK PAGE

- 6 Soluble salts can be made using a base and an acid.
 - (a) Complete this method of preparing dry crystals of the soluble salt cobalt(II) chloride-6-water from the insoluble base cobalt(II) carbonate.

step 1 Add an excess of cobalt(II) carbonate to hot dilute hydrochloric acid.

(b) (i) 5.95g of cobalt(II) carbonate were added to 40 cm^3 of hydrochloric acid, concentration 2.0 mol/dm^3 .

Calculate the maximum yield of cobalt(II) chloride-6-water and show that the cobalt(II) carbonate was in excess.

 $CoCO_3 \ + \ 2HC\mathit{l} \ \rightarrow \ CoC\mathit{l}_2 \ + \ CO_2 \ + \ H_2O$

 $CoCl_2$ + $6H_2O \rightarrow CoCl_2.6H_2O$

maximum yield:

	number of moles of HCl used =
	number of moles of CoCl ₂ formed =
	number of moles of CoCl ₂ .6H ₂ O formed =
	mass of one mole of $CoCl_2.6H_2O = 238g$
	maximum yield of CoCl ₂ .6H ₂ O =g
	to show that cobalt(II) carbonate is in excess:
	number of moles of HC <i>l</i> used = (use your value from above)
	mass of one mole of $CoCO_3 = 119g$
	number of moles of $CoCO_3$ in 5.95g of cobalt(II) carbonate =[5]
(ii)	Explain how these calculations show that cobalt(II) carbonate is in excess.
	[1]
	[Total: 10]

7 Iodine reacts with chlorine to form dark brown iodine monochloride.

$$I_2 \ + \ C\mathit{l}_2 \ \rightarrow \ 2IC\mathit{l}$$

This reacts with more chlorine to give yellow iodine trichloride. An equilibrium forms between these iodine chlorides.

(a) What do you understand by the term *equilibrium*?

[2]

(b) When the equilibrium mixture is heated, it becomes a darker brown colour. Suggest if the reverse reaction is endothermic or exothermic. Give a reason for your choice.

[1]

(c) The pressure on the equilibrium mixture is decreased.

(i) How would this affect the position of equilibrium? Give a reason for your choice.

	It would move to the		
	reason		
			[1]
(ii)	Describe what you wo	uld observe.	

[1]

(d) Calculate the overall energy change for the reaction between iodine and chlorine using the bond energy values shown.

$$I_2 + Cl_2 \rightarrow 2ICl$$

Bond	Energy / kJ per mol
I–I	151
Cl–Cl	242
I–Cl	208

Show your working.

[3]

(e) Draw a labelled energy level diagram for the reaction between iodine and chlorine using the information in (d).

[2]

[Total: 10]

- 8 The alcohols form an homologous series.
 - (a) Give three characteristics of an homologous series.

[3]

(b) The following two alcohols are members of an homologous series and they are isomers.


 $CH_3 - CH_2 - CH_2 - CH_2 - OH$ and $(CH_3)_2CH - CH_2 - OH$

(i) Explain why they are isomers.

[2]

(ii) Deduce the structural formula of another alcohol which is also an isomer of these alcohols.


[1]




(d) The alcohol ethanol can be made by fermentation. Yeast is added to aqueous glucose.

 $C_6H_{12}O_6(aq) \rightarrow 2C_2H_5OH(aq) + 2CO_2(g)$

Carbon dioxide is given off and the mixture becomes warm, as the reaction is exothermic. The graph shows how the rate of reaction varies over several days.



(i) Suggest a method of measuring the rate of this reaction.



BLANK PAGE

- **9** There are two types of polymerisation, addition and condensation.
 - (a) Explain the difference between these two types of polymerisation.

[2]

(b) Some plastics, formed by polymerisation, are non-biodegradable.

Describe two pollution problems that are caused by non-biodegradable plastics.

[2]

(c) The polymer known as PVA is used in paints and adhesives. Its structural formula is shown below.

--CH₂--CH---CH₂--CH---| | | OCOCH₃ OCOCH₃

Deduce the structural formula of its monomer.

(d) A condensation polymer can be made from the following monomers.

HOOC(CH₂)₄COOH and H₂N(CH₂)₆NH₂

Draw the structural formula of this polymer.

[3]

[Total: 8]

								Group	dni								
-	=											=	≥	>	5	-IIV	1
							۲										2
							т										θH
				Key			hydrogum 1										helium 4
	4		ñ	atomia number	e.							6		2			10
-	Be		ato	atomic symbol	poq							в	υ	z	0	u.	Ne
lithium	beryflum			TRUTTED.								liorun	carbon	mèrogen	cas) gan	fluction	1000
~	a		relati	relative atomic mass	mass							1	12	14	16	19	50
1	12											13	4	16	16	17	18
Ra	BM											Ĩ	ō	۵.	s	õ	Ł
modum 23	nagrasium 24											ahminum 27	allican 28	phosphonus 31	auftur 32	dhlorine 35.5	argon 40
6	8	21	22	33	54	25	8	27	28	29	8	31	32	8	ş	35	98
×	Сa	ŝ	F	>	ບັ	Mn	e L	ů	ï	5	Zn	Ga	ge	As	Se	'n	고
potessium	caloum	scandium	13 ar kum	vanadium	chromium	manganese	iron	odult	rickel	Jodebo	2010	gallum	gemanium	amenic	selenium	bromine	krypton
8	6	45	48	5	25	55	8	88	59	8	8	۶	73	22	62	8	\$
37	38	8	ş	4	4	43	\$	\$	\$	4	\$	49	3	5	23	53	3
ß	ທັ	≻	Ż	đ	еŴ	ř	2	문	P	Ag	8	Ч	Sn	å	Te	I	Xe
ndichun	atrantium ac	ytinum 20	zinconum 24	riothum 200	motyboinum	technolium	nuhanium 404	modum 2004	pulsdium 400	allour 4.00	audmium 4.4.0	indium A 4 4	÷ ;	artimory	to built m	indire 4.07	201CU
8 5	00	R0	R	26	8 7	1 20	5 8	3	901	<u>60</u>	711	110	8	771	07	171	131
8 6	8 2	D//1 forthandits	71	2	2 3	2	۹ (Ξ,	2	2	8 -	ōŀ	5 6	2 2	s é	8 3	8 6
3	e Ca		Ē	m -	2	Ŷ	ŝ	5	ĩ	PA -	ĝ	2	2	'n	9 2	¥	ž
133	137		178	181	184	186 186	190	190 190	195	2006 197	201	204	207	209	muuood	esione	10000
87	88	89-103	104	105	106	107	108	100		111	112		114		116		
Ŀ	Ra	actinoids	ž	90	ß	旧	Чŝ	Mt	S	Ra	ວົ		F١		2		
fisroium	rsoum		ntherfordism	addrium	setborgkun	bohrůum	hassium	meltrenum	5	roenigenium	openisium		flerovium		Rvemorium		
	I		1				I	1	I	1					1		
		25	58	65	8	61	62		64		99	67	68	8	70	14	
lanthanoids	ds	La	ő	ŗ		Б	SB		В		2	위	ա	Ta	Υb	3	
		lar/hanum	cerium	prano dimiure	Ę	promothum	samarium	europium	gadolinium		dysprosium	holmàum	ettium	thulium	yterbium	h/relium	
		139	140	141	144	I	150		157		163	165	167	169	173	175	
		8	90	9	92	8	8		96		96	8	100	101	102	8	
actinoids		Ac	£	Ра	⊐	ď	Ъ	Am	ő	ň	õ	ů	Ē	ΡW	٩	5	
		actinium	therium	protectinium	uranium	neptunium	ph/confurm	americium	ourium	terkeitum	californium	oinstehrium	femium	merclelovium	nobelium	lawrondum	
		ı	232	231	238	1	1	'	1	1	I	'	1	'	ı	ı	
The volume of one mole of any case is 24 dm ³ at room temperature and pressure (r.t.p.)	ne of on	a mole of	anv dae	ie 24 dm ³	at more	homorah	n han ord	Neering									

ŝ × . 5

Permission to reproduce items where third-party owned material protected by copyright is included has been sought and cleared where possible. Every reasonable effort has been made by the publisher (UCLES) to trace copyright holders, but if any items requiring clearance have unwittingly been included, the publisher will be pleased to make amends at the earliest possible opportunity.

Cambridge Assessment International Education is part of the Cambridge Assessment Group. Cambridge Assessment is the brand name of University of Cambridge Local Examinations Syndicate (UCLES), which is itself a department of the University of Cambridge.